

Cal/Ecotox

Toxicity Data for Loggerhead Shrike (*Lanius ludovicianus*)*

Page 1

Chemical	Tox Exposure	Endpoint Type	Endpoint Description	Endpoint Value	Note	Reference
AMMONIUM NITRATE	54 kg/ha application rate	TOX-Non-Repro-Sublethal - indirect effects	loss of eggs, nestlings, fledglings and adults post-spray vs. unsprayed controls	increased	a	1
AMMONIUM NITRATE	54 kg/ha application rate	TOX-Non-Repro-Sublethal - indirect effects	percent increase in breeding territory size post-spray vs. unsprayed controls	significantly increased (138.5% +/- 86.1 SD)	b	1
AMMONIUM NITRATE	54 kg/ha application rate	TOX-Non-Repro-Sublethal - indirect effects	timing of nesting season termination in pairs post-spray vs. unsprayed controls	significantly earlier	c	1
DDD (4,4'-); DDE (4,4'-); DDT (4,4'-)	0.22 kg DDT/ha	TOX-EXP IND - accumulation	concentrations of DDT, DDE, DDD (combined) in skin and brains, 2 years post-application	293.43-820.93 ppm, lipid weight	d	2
DDD (4,4'-); DDE (4,4'-); DDT (4,4'-)	0.22 kg DDT/ha	TOX-EXP IND - accumulation	figure showing concentration of DDT and metabolites in skin and brains before and after application	see figure	e	2
DDE (4,4'-)	21.04 (3.51 SE) ppm in fat of adult females	TOX-EXP IND - accumulation	concentration in eggs	3.09 (0.09 SE) ppm, wet wt	f	3
DDE (4,4'-)	21.04 (3.51 SE) ppm in fat of adult females	TOX-REPRO - physiology	eggshell thickness index compared with reference samples collected in 1875-95	decreased	g	3
DDE (4,4'-)	21.04 (3.51 SE) ppm in fat of adult females	TOX-REPRO - physiology	linear negative correlation of egg DDE concentration and shell thickness	y = 92.610 - 2.412 log x	h	3
DIELDRIN	0, 1, 2, 4, 8 mg/kg bw/d (87% active ingredient)	TOX-MORT - dose-response data	rate of mortality compared with controls	increased at 1, 2, 4, 8 mg/kg/d	i	4
DIELDRIN	0, 1 mg/kg bw/d (87% active ingredient)	TOX-Non-Repro-Sublethal - behavioral effects	latency to formation of natural prey killing behaviors	increased at 1 mg/kg	j	4

Notes

- a Both Adult and Juv.; FL; B; Species - California (R)=*Lanius ludovicianus*; TOX - Chemical=6484-52-2; N=4 control, 8 treatment territories; June-July; MacArthur Agroecology Research Center, Archbold Biological Station; Tox Exp Tech=application to habitat; Tox Exp Dur=approx. 1 month; Tox Study Dur=approx. 1 month; Tox Stat Sig=NR
- b Both Adult and Juv.; FL; B; Species - California (R)=*Lanius ludovicianus*; TOX - Chemical=6484-52-2; N=4 control, 8 treatment territories; June-July; MacArthur Agroecology Research Center, Archbold Biological Station; Tox Exp Tech=application to habitat; Tox Exp Dur=approx. 1 month; Tox Study Dur=approx. 1 month; Tox Stat Sig=Y; Total number of insects and insect species collected in sweepnets two weeks post-spray was lower than that in unsprayed pasture.
- c Both Adult and Juv.; FL; B; Species - California (R)=*Lanius ludovicianus*; TOX - Chemical=6484-52-2; N=4 control, 8 treatment territories; June-July; MacArthur Agroecology Research Center, Archbold Biological Station; Tox Exp Tech=application to habitat; Tox Exp Dur=approx. 1 month; Tox Study Dur=approx. 1 month; Tox Stat Sig=Y
- d NR; Yolo; CA; NR; Species - California (R)=*Lanius ludovicianus*; TOX - Chemical=72-54-8; TOX - Chemical=72-55-9; TOX - Chemical=50-29-3; N=2 birds; 3.2 km NE of Davis; Tox Exp Tech=experimental application; Tox Exp Dur=1 application; Tox Study Dur=2 yrs; Tox Stat Sig=NR
- e NR; Yolo; CA; NR; Species - California (R)=*Lanius ludovicianus*; TOX - Chemical=72-54-8; TOX - Chemical=72-55-9; TOX - Chemical=50-29-3; N=27 birds (3-4/sampling time); 3.2 km NE of Davis; Tox Exp Tech=experimental application; Tox Exp Dur=1 application; Tox Study Dur=2 yrs; Tox Stat Sig=NR; See citation for residues in soil, vegetation, invertebrates, and mammals over time after application.
- f Embryo; IL; NR; Species - California (R)=*Lanius ludovicianus*; TOX - Chemical=72-55-9; N=20 clutches, 104 eggs; Apr, May; southern IL; Tox Exp Dur=NR; Tox Exp Dur=parental transfer; Tox Study Dur=NR; Tox Stat Sig=NR; Egg and female shrike samples were collected independently, but within same geographic location.
- g Embryo; IL; NR; Species - California (R)=*Lanius ludovicianus*; TOX - Chemical=72-55-9; N=20 clutches, 104 eggs; Apr, May; southern IL; Tox Exp Dur=parental transfer; Tox Exp Dur=NR; Tox Study Dur=NR; Tox Stat Sig=Y
- h Embryo; IL; NR; Species - California (R)=*Lanius ludovicianus*; TOX - Chemical=72-55-9; N=20 clutches, 104 eggs; Apr, May; southern IL; Tox Exp Dur=parental transfer; Tox Exp Dur=NR; Tox Study Dur=NR; Tox Stat Sig=Y
- i Juvenile; Lab; NR; Species - California (R)=*Lanius ludovicianus*; TOX - Chemical=60-57-1; TOX - Dose-Response Data Format=DR Table; N=10-11/dose; Age=1 day post-hatch (begin); collected in San Diego County, CA; Tox Exp Tech=gavage; Tox Exp Dur=until death (max. 103 d); Tox Study Dur=until death (max. 103 d); Tox Stat Sig=NR; See table in citation for mean ages at death for different doses (range 16.8 d @ 8 mg/kg/d to 84.5 d @ 1 mg/kg/d)
- j Juvenile; Lab; NR; Species - California (R)=*Lanius ludovicianus*; TOX - Chemical=60-57-1; TOX - Dose-Response Data Format=DR Figure; TOX - Dose-Response Data Format=DR Table; N=10-11/dose; Age=1 day post-hatch (begin); collected in San Diego County, CA; Tox Exp Tech=gavage; Tox Exp Dur=until death (max. 103 d); Tox Study Dur=>55 d (behavioral testing); Tox Stat Sig=Y; See citation for table and figures showing timing of occurrence of various mouse killing behaviors.

References

- 1 Yosef, Reuven, and Mark A. Deyrup. 1998. Effects of fertilizer-induced reduction of invertebrates on reproductive success of loggerhead shrikes (*Lanius ludovicianus*). J. Zool. 139:307-312.

2 Rudd, R.L., R.B. Craig and W.S. Williams. 1981. Trophic accumulation of DDT in a terrestrial food web. Environ. Pollut. Ser. A. 25:219-228.
3 Anderson, William L., and Ronald E. Duzan. 1978. DDE residues and eggshell thinning in loggerhead shrikes. Wilson Bull. 90(2):215-220.
4 Busbee, Everette L. 1977. The effects of dieldrin on the behavior of young loggerhead shrikes. Auk. 94:28-35.

* Cal/EPA, OEHHA and the University of California Regents are not responsible for damages of any kind resulting from the use of or reliance on information in this report. Users are encouraged to consult the original data. Updated: February 1999.